

Weak Convergence Methods for Approximation of Evaluation of Path-dependent Functionals*

Qingshuo Song,[†]G. Yin,[‡]Qing Zhang[§]

February 19, 2013

Abstract

This paper develops approximation methods for path-dependent functionals, which have been used in many applications involving path-dependent objective functions. By a first glance, the problem may appear as a standard approximation of a stopping time problem using traditional techniques. Nevertheless, a closer scrutiny would reveal that the problem under consideration is far more challenging and difficult. The difficulty is mainly because the traditional dynamic programming approach falls apart, not to mention any hope for a closed-form solution or any viable numerical PDE methods. This paper focuses on approximation methods in Monte Carlo simulations. To the best of our knowledge, convergence analysis is available only in few special cases due to the complexity of the nature of path dependence. In contrast to the traditional approach, this work provides a non-traditional convergence method in Monte Carlo analysis, and establishes a general framework for the convergence analysis using the Markov chain approximations. In particular, the approach is based on actual computations under the Skorohod topology. Some examples such as the approximation of discretely monitoring barrier option are considered.

*This research was supported in part by the Research Grants Council of Hong Kong No. CityU 103310, and in part by in part by the National Science Foundation under DMS-1207667.

[†]Department of Mathematics, City University of Hong Kong. 83 Tat Chee Avenue, Kowloon Tong, Hong Kong, song.qingshuo@cityu.edu.hk.

[‡]Department of Mathematics, Wayne State University, Detroit, Michigan 48202, gyin@math.wayne.edu.

[§]Department of Mathematics, The University of Georgia, Athens, GA 30602, qingz@math.uga.edu.

1 Introduction and Examples

In many applications, one needs to evaluate path-dependent objective functions. Many of such calculations require the handling of stopping times in which the stopping boundaries are non-constant but are path dependent. The applications stem from evaluations of functionals in networked systems as well as in finance. In addition, they also arise in Euler's approximation to solution of stochastic differential equations. This paper is concerned with approximation methods for computing such objective functions. By a first glance, the problem may appear as a standard approximation of a stopping time problem involving traditional techniques. Nevertheless, a closer scrutiny would reveal that the problem under consideration is far more challenging and difficult. Even though the problem may still be viewed as stopping time problems, the targets for the stopping times are not fixed constants as most of those treated in the literature, but rather a path-dependent criteria. This is where the main difficulty arises because the traditional dynamic programming and Markovian properties fall apart.

To begin, there are virtually no closed-form solutions for the path-dependent objective functionals. One has to look for alternatives. The next possibility is numerical approximation using partial differential equation (PDE) based techniques. Nevertheless, such possibility is ruled out due to the lack of Markovian properties. The only often used technique left is the Monte Carlo method. While most of the existing methods for treating Monte Carlo procedures are somewhat ad hoc, this work develops a systematic alternative method for analyzing the convergence of the approximation algorithm. In this paper, we use a weak convergence approach and carry out the convergence analysis under the Skorohod topology. One of the main ingredients is the use of a generalized projection operator. With the use of such projections, we proceed to by exploring the intrinsic properties of the Skorohod space. The convergence analysis developed in this paper provides a thorough understanding of the nature of path dependence and a general framework for handling many path-dependent problems.

1.1 Path-dependent Objective Functions

We are interested in approximating path-dependent functions in a finite-time horizon. For simplicity, we focus our attention on the time interval $[0, 1]$ to avoid using more complex notation. Let $C[0, 1]$ be the collection of continuous real-valued functions defined on $[0, 1]$, and $D[0, 1]$ be the collection of all RCLL (right continuous with left-hand limits) functions on $[0, 1]$. With $\mathbb{F} = \{\mathcal{F}_t : t \in [0, 1]\}$ and $\mathcal{F} = \mathcal{F}_1$, let $(\Omega, \mathcal{F}, \mathbb{P}, \mathbb{F})$ be a filtered probability space satisfying the usual conditions; see [10]. As a convention, we use capital letters to denote random elements and use lower case letters for deterministic (or non-random) elements. For instance, $x \in D[0, 1]$ is a

RCLL path defined on $[0, 1]$, while $X : \Omega \mapsto D[0, 1]$ is a random RCLL process with sample paths in $D[0, 1]$. As usual, X can be regarded as a function of two variables time t and sample point ω . That is, for each fixed $\omega \in \Omega$, $X(\cdot, \omega)$ denotes a sample path and for each fixed t , $X(t, \cdot)$ is a random element in $\mathbb{R}$.

Throughout the paper, $\alpha, \beta \in C[0, 1]$ are given satisfying $\alpha(t) < \beta(t)$ for all $t \in [0, 1]$. The domain of our interest is defined by all the points bounded by α and β , that is,

$$Q = \{(a, t) : \alpha(t) < a < \beta(t), t \in [0, 1]\}.$$

The cross-section $Q(t) = (\alpha(t), \beta(t)) \subset \mathbb{R}$ may be time-dependent. We also denote its boundary by

$$\partial Q = \{(\alpha(t), t) : t \in [0, 1]\} \cup \{(\beta(t), t) : t \in [0, 1]\}.$$

We study an $\mathbb{F}$ -adapted continuous process $X : [0, 1] \times \Omega \mapsto C[0, 1]$. Let τ be the first hitting time to the boundary of the time-dependence domain

$$\tau = \inf\{t > 0 : X(t) \notin Q(t)\} \wedge 1. \quad (1)$$

Given a positive integer m , let $(c_1, c_2, \dots, c_m)' \in [0, 1]^m$ be a column vector, representing a vector of scaling factors. Define a set of random times

$$\mathcal{T} = \{\tau_j = c_j \tau : 1 \leq j \leq m\}.$$

Note that random times τ_j are measurable with respect to $\mathcal{F}_1$, but may not be a stopping time whenever $0 < c_j < 1$. Now we introduce the notion of generalized projection operator by defining Π as

$$\Pi_{\mathcal{T}}(X) = \{X(\tau_j) : 1 \leq j \leq m\} \in \mathbb{R}^m.$$

The Π is a generalized projection in the sense that it distinguishes from the conventional projection of a path by projecting X to a collection of random times instead of fixed times; see [3].

For $x \in D[0, 1]$, we define a maximum process x^* by $x^*(t) = \sup_{0 \leq s \leq t} x(s)$. It is easy to check that $x^* \in D[0, 1]$. For some measurable function $g : \mathbb{R}^{m+2} \rightarrow \mathbb{R}$, we are interested in the computation of the objective functional $V : C \mapsto \mathbb{R}$ defined by

$$V = \mathbb{E}[g(\Pi_{\mathcal{T}}(X), X^*(\tau), \tau)]. \quad (2)$$

The value function may be a measurable function of the initial states $V(x, t)$ if the functional V is given as a non-path-dependent form of $\sup \mathbb{E}[f(X(T))]$ for some function $f : \mathbb{R} \mapsto \mathbb{R}$. However, we emphasize that our main interest is the computation when the functional is path-dependent in the form of (2), which has a wide range of applications in mathematical finance.

Remark 1 Related literature in connection with a hitting time under a Markovian framework can be found in Dufour and Piunovskiy [6], de Saporta, et al. [5], and Szpruch and Higham [18] among others. In particular, in [6], the existence of optimal stopping with constraints is established using a convex analytic approach. A numerical method for optimal stopping of piecewise deterministic Markov processes is considered in [5]. Bounds for the convergence rate of their algorithm are obtained by introducing quantization of the post jump location and path-adapted time discretization grids. In [18], an application in physics (thermodynamic limit) involving mean hitting time behavior is considered.

1.2 Examples

The evaluation of path-dependent objective functions is of great interest in many networked systems as well as in financial applications such as the option pricing. To illustrate, consider the underlying stock price X that is a non-negative martingale process under $\mathbb{P}$ that is the corresponding *equivalent local martingale measure*. Assuming zero interest rate, then the value V of (2) can be considered as a general form of a class of option prices including look-back option, rebate/barrier option, Asian option, Bermuda option etc. For illustration purpose, let us fix

$$\mathcal{T} = \left\{ \frac{i}{m}\tau : 1 \leq i \leq m \right\}.$$

1. (Barrier option) Let $-\alpha(t) = \infty$ and $\beta(t) = 1$, and thus $\tau = \inf\{s > t : X(s) \notin (-\infty, 1)\} \wedge 1$. The price formula of the up-and-in barrier call with strike 1/2 at maturity $T = 1$ is given by

$$V = \mathbb{E}[(X(1) - \frac{1}{2})^+ I_{\{\tau \leq 1\}}].$$

In this case, the payoff function is

$$g(x_1, x_2, \dots, x_m, x_{m+1}, t) = (x_m - \frac{1}{2})^+ I_{\{t \leq 1\}} = (x_m - \frac{1}{2})^+ I_{\{x_{m+1} \geq 1\}},$$

which is discontinuous.

2. (Discretely monitoring barrier option) As mentioned in [4] and [12], treating a class of so-called monitoring barrier options, most models assume continuous monitoring of the barrier like the above. However in practice most, if not all, barrier options traded in markets are discretely monitored. Unlike their continuous counterparts, there is essentially no closed form solution available, and even numerical pricing is difficult. For instance, again we set the same barriers $-\alpha(t) = \infty$ and $\beta(t) = 1$. However, we will replace τ in the above continuous time barrier option by

$$\hat{\tau} = \inf\{s = i/m : X(s) \notin (-\infty, 1), i = 1, 2, \dots, m\} \wedge 1,$$

The price formula of up-and-in barrier call with strike $\frac{1}{2}$ is then

$$V = \mathbb{E}\left[\left(X(1) - \frac{1}{2}\right)^+ I_{\{\hat{\tau} \leq 1\}}\right]. \quad (3)$$

Observe that V can be rewritten as $V = \mathbb{E}[(X(1) - \frac{1}{2})^+ I_{\{\max(\Pi_{\mathcal{T}}(X)) \geq 1\}}]$, and the payoff function g is accordingly

$$g(x_1, x_2, \dots, x_m, x_{m+1}, t) = \left(x_m - \frac{1}{2}\right)^+ I_{\{\max\{x_1, \dots, x_m\} \geq 1\}}.$$

Note that g has linear growth, is unbounded, and is discontinuous.

1.3 Computational Methods

To evaluate the path-dependent objective functions, a major difficulty arises from the lack of Markovian conditions. This rules out the possibility of any PDE based methods including the finite difference or finite element methods. In addition, the time-dependent barriers α and β also create additional layer of difficulty. In this paper, we consider Monte Carlo methods to obtain a feasible estimation of the path-dependent problems. Monte Carlo method is a class of computational algorithms that relies on repeated random sampling to evaluate its deterministic value using its probabilistic fact. This includes Euler-Maruyama approximation [11] and Markov chain approximation [8], [13], and [14] among others.

The general idea of the Monte Carlo method in this vein is the following. For each sample point $\omega \in \Omega$, use $X^h(\cdot, \omega) \in D[0, 1]$ to denote a simulated path for the underlying process $X(\cdot, \omega) \in C[0, 1]$ by a certain Monte Carlo method with a small parameter h . Define the approximated stopping time and projection operator by

$$\tau^h = \inf\{t > 0 : X^h(t) \notin (\alpha(t), \beta(t))\} \wedge 1 \text{ and } \mathcal{T}^h = \{\tau_j^h = c_j \tau^h : 1 \leq j \leq m\}.$$

One can approximate V in this way by computing

$$V^h = \mathbb{E}[g(\Pi_{\mathcal{T}^h}(X^h), X^{h,*}(\tau^h), \tau^h)], \quad (4)$$

where

$$\Pi_{\mathcal{T}^h}(X^h) = \{X^h(\tau_j^h) : 1 \leq j \leq m\}.$$

The goal is to design an appropriate Monte Carlo method so that the desired convergence takes place eventually, i.e.,

$$\lim_{h \rightarrow 0} V^h = V.$$

As the objective value is given as an expectation form of (2), V is invariant under the same distribution, and the usual requirements for the Monte Carlo method are intuitively given as follows:

(H1) X^h converges in distribution to X as $h \rightarrow 0$, denoted by $X^h \Rightarrow X$.

(H2) g is continuous.

A couple of natural questions are as follows.

- Are (H1)-(H2) sufficient to guarantee the desired convergence $\lim_{h \rightarrow 0} V^h = V$?
- Can (H2) be possibly weakened to some discontinuous function g so that the barrier option pricing (see Section 1.2) can be included?

1.4 Counter Examples

Interestingly, there exist circumstances that lead to counter examples in connection with the desired convergence under (H1)-(H2).




Figure 1: Demonstration of tangency problem: The sample paths of the two processes (solid line and dotted line) are fairly close to each other, but their first exit times are far apart indicating a loss of continuity.

A noticeable counter example is given in Figure 1, which is motivated by the so-called *tangency problem*. Consider two underlying processes X^1 (solid line) and X^2 (dotted line) in Figure 1. No matter how close X^1 and X^2 are at the initial states, the difference between their first exit time τ_1 and τ_2 could be very far away; see [2]. The above idea is formulated as the following example.

Example 1 Let $\{X(s) : 0 \leq s \leq 1\}$ be a deterministic process given by $X(s) = 1 - (s - \frac{1}{2})^2$. Let $\alpha(s) = -\infty$ and $\beta_s = 1$ for all $s \geq 0$. Then the exit time of X is

$$\tau = \inf\{s > 0 : X(s) \notin (\alpha(s), \beta(s))\} \wedge 1 = \frac{1}{2}.$$

Define a family of the processes parameterized by h with $X^h(s) = X(s) - h$. Although X^h converge to X in L^∞ as $h \rightarrow 0^+$, we note that

$$\tau^h = \inf\{s > 0 : X^h(s) \notin (\alpha(s), \beta(s))\} \wedge 1 = 1$$

not converging to $\tau = 1/2$.

The next example is different from the first one in the sense that there is no obstacles and the underlying process is taken from a Bessel process. This system is often used in finance to model the dynamics of asset prices, of the spot rate and of the stochastic volatility, or as a computational tool. In particular, computations for the celebrated Cox-Ingersoll-Ross (CIR) and Constant Elasticity Variance (CEV) models can be carried out using Bessel processes.

Example 2 Let $X(t)$ be the Bessel process of dimension 3 with initial $X(0) = 1$, i.e.,

$$X(t) = 1 + W(t) + \int_0^t \frac{ds}{X(s)}, \quad (5)$$

which is a well known strict local martingale with $\mathbb{E}[X(1)] < 1$; see Page 336 of [9]. Define $X^h(t) = X(t) \wedge \frac{1}{h}$. Fix an arbitrary partition $\mathcal{T}_n = \{t_1 \leq \dots \leq t_n\} \subset [0, 1]$, and an open set $O \subset [0, N]^n$. Note that, for any $h \in (0, 1/N)$, we have

$$\mathbb{P}\{\Pi_{\mathcal{T}_n}(X^h) \in O\} = \mathbb{P}\{\Pi_{\mathcal{T}_n}(X) \in O\} - \mathbb{P}\{\Pi_{\mathcal{T}_n}(X) \in O, \Pi_{\mathcal{T}_n}(X^*) \geq \frac{1}{h}\}$$

and

$$\begin{aligned} 0 &\leq \mathbb{P}\{\Pi_{\mathcal{T}_n}(X) \in O, \Pi_{\mathcal{T}_n}(X^*) \geq \frac{1}{h}\} \\ &\leq \mathbb{P}\{\Pi_{\mathcal{T}_n}(X^*) \geq \frac{1}{h}\} \rightarrow 0 \quad \text{as } h \rightarrow 0. \end{aligned}$$

This implies $\lim_{h \rightarrow 0} \mathbb{P}\{\Pi_{\mathcal{T}_n}(X^h) \in O\} = \mathbb{P}\{\Pi_{\mathcal{T}_n}(X) \in O\}$, and thus $\Pi_{\mathcal{T}_n}(X^h) \Rightarrow \Pi_{\mathcal{T}_n}(X)$ due to the arbitrariness of the set O . Moreover, Kolmogorov consistency theorem together with the uniqueness of the weak solution of SDE (5) implies that $X^h \Rightarrow X$ due to the arbitrariness of n and $\mathcal{T}_n$. However, since X^h is a bounded local martingale, hence a martingale, we have the following undesired convergence: $\lim_{h \rightarrow 0} \mathbb{E}[X^h(1)] = 1 > \mathbb{E}[X(1)]$.

1.5 Goal and Outline of the Paper

The above examples show that (H1)-(H2) may not be sufficient for Monte Carlo simulation to be convergent to the right value of (2). This propose the following question:

(Q1) Given $X^h \Rightarrow X$, what is a sufficient condition for the convergence $\lim_{h \rightarrow 0} V^h = V$? Is it possible to weaken the continuity of g to cover the barrier option?

In this paper, we first explore a rigorous proof of the convergence in Section 2. The results in Theorem 6 provide a set of sufficient conditions for the convergence. It can also explain why Examples 1 and 2 do not work.

Our approach is based on actual computations using the Skorohod metric in the $D[0, 1]$ space. This is the first attempt in this context to the best of our knowledge. Such an approach is advantageous. For example, different types of options mentioned the convergence of Monte Carlo methods have been studied separately; see [8] and the references therein. However, Theorem 6 provides a unified theoretical basis for the convergence in a general path-dependent form with possibly discontinuous payoff function g and finite time-dependent barrier $(\alpha(t), \beta(t))$. This covers a wider range of applications of Theorem 6 to various types of options, for instance, barrier options, Asian options, look back options, etc. It is also notable that the assumptions on the barrier in the path-dependent problem has some similarities to the assumptions imposed for stochastic control exit problems; see [2]. To complete the approximation of the value V of (2), one should consider the following question in addition to (Q1),

(Q2) Given X , how does one construct approximating processes X^h such that $X^h \Rightarrow X$?

Among many of the possible answers to (Q2), we will mainly focus on the construction of X^h in the framework of Markov chain approximation in Section 3. Recall that from the book [14], a family of continuous approximating processes X^h constructed using Markov chain is convergent to X in distribution if the Markov chain satisfies the local consistency [14, equation (9.4.2)]; see also [14, Theorem 10.4.1]. Compared to [14], we investigate the weak convergence for more generalized local consistency condition. This enables us to cover various types of Monte Carlo simulations in the framework of Markov chain approximation to verify its weak convergence using the generalized local consistency. It essentially extends the use of Markov chain approximation.

To proceed, the rest of the paper is arranged as follows. Section 2 provides sufficient conditions for convergence of our procedure. Then the convergence of the algorithms is established. Section 3 presents the weak convergence of Markov chain approximation under the generalized local consistency. Also provided here is the answer: Can we get strong convergence? Finally, the paper is concluded with some further remarks in Section 4.

2 Sufficient Conditions for Convergence

2.1 Preliminaries

We use the notation given in [3]. Define a metric on $D[0, 1]$ by

$$\|x - y\| = \sup_{t \in [0, 1]} |x(t) - y(t)|, \quad \forall x, y \in D[0, 1].$$

Then, the continuous function space on $[0, 1]$, denoted by $C[0, 1]$, is complete with respect to the uniform topology with the above metric $\|\cdot\|$. On the other hand, the RCLL function space defined on $[0, 1]$, denoted by $D[0, 1]$, is equipped with the Skorohod topology with the metric

$$\|x - y\|_s = \inf_{\lambda \in \Lambda} \{\|\lambda - I\|, \|x \circ \lambda - y\|\}, \quad \forall x, y \in D[0, 1],$$

where $x \circ y$ denotes the composite function of x and y , and Λ is the collection of all continuous increasing functions λ on $[0, 1]$ with $\lambda(0) = 0$ and $\lambda(1) = 1$. Note that $D[0, 1]$ is not complete under the metric $\|\cdot\|_s$, but it is complete under an equivalent metric $\|\cdot\|_s^o$ defined by

$$\|x - y\|_s^o = \inf_{\lambda \in \Lambda} \left\{ \sup_{s < t} \left| \log \frac{\lambda(t) - \lambda(s)}{t - s} \right|, \|x - y\| \right\}, \quad \forall x, y \in D[0, 1].$$

For notational convenience, we will adopt $\|\cdot\|_s$ in the rest of the paper. Recall that, for $x \in D[0, 1]$, we define $x^* \in D[0, 1]$ by $x^*(t) = \sup_{0 \leq s \leq t} x(s)$.

Lemma 2 *Let $\pi^* : D[0, 1] \mapsto D[0, 1]$ by $\pi^*(x) = x^*$. Then, π^* is continuous with respect to the Skorohod topology.*

PROOF: Since λ is strictly monotone, $x^* \circ \lambda = (x \circ \lambda)^*$. Therefore,

$$\begin{aligned} \|x^* - y^*\|_s &= \inf_{\lambda \in \Lambda} \{ \|\lambda - I\|, \sup_{t \in [0, 1]} |x^*(\lambda(t)) - y^*(t)| \} \\ &= \inf_{\lambda \in \Lambda} \{ \|\lambda - I\|, \sup_{t \in [0, 1]} |(x \circ \lambda)^*(t) - y^*(t)| \} \\ &= \inf_{\lambda \in \Lambda} \{ \|\lambda - I\|, \sup_{t \in [0, 1]} \|x \circ \lambda - y\|_{[0, t]} \} \\ &= \inf_{\lambda \in \Lambda} \{ \|\lambda - I\|, \|x \circ \lambda - y\| \} = \|x - y\|_s. \end{aligned}$$

So, π^* is continuous in $D[0, 1]$ with respect to the Skorohod topology. $\square$

2.2 Main Convergence Results

We make the following assumptions.

(A1) If $\beta < \infty$, then $\mathbb{P}\{(X - \beta)^*(1) \geq 0\} > 0$. If $\alpha > -\infty$, then $\mathbb{P}\{(-X + \alpha)^*(1) \geq 0\} > 0$.

Note that Condition (A1) means that the process hits both barriers α and β with positive probability, whenever $-\alpha$ or β are bounded above. This is not a restriction, since one can simply set $\alpha(t) = -\infty$ (resp., $\beta(t) = \infty$) for all $t \in [0, 1]$, if X never hits α (resp., β) almost surely in $\mathbb{P}$. If $-\alpha(t) = \beta(t) = \infty$, then $\tau = 1$.

(A2) X satisfies $\inf\{t > \tau : X(t) \notin [\alpha(t), \beta(t)]\} = \tau$ almost surely $\mathbb{P}$.

Condition (A2) requires that the boundary ∂Q is regular with respect to the process X . Note that, for any small $\varepsilon > 0$, X exits from $\bar{Q}$ in the interval $(\tau, \tau + \varepsilon)$ under (A2). Loosely speaking, the condition (A2) means that the process $X(t)$ exits $\bar{Q}$ immediately after it hits the boundary at τ . More discussions are referred to Remark 7.




Figure 2: Illustration for the partition of $C[0, 1]$

To proceed, let's partition the space $C[0, 1]$ as follows: Define

$$C_1 = \{x \in C[0, 1] : \pi(x) < 1, x(\pi(x)) = \beta(\pi(x)), \inf\{t > \pi(x) : x(t) > \beta(t)\} = \pi(x)\},$$

$$C_2 = \{x \in C[0, 1] : \pi(x) < 1, x(\pi(x)) = \alpha(\pi(x)), \inf\{t > \pi(x) : x(t) < \alpha(t)\} = \pi(x)\},$$

and

$$C_3 = \{x \in C[0, 1] : \pi(x) = 1\}, \text{ and } C_4 = C[0, 1] \setminus (\cup_{i=1}^3 C_i).$$

Accordingly, we can write $C[0, 1] = \cup_{i=1}^4 C_i$ and $C_i \cap C_j = \emptyset$ for $i \neq j$. As for the illustration, one can see that the four curves depicted in Figure 2 belong to four different subsets separately, that is, $l_i \in C_i$ for $i = 1, 2, 3, 4$. Note that (A2) implies that

$$\mathbb{P}\{X \in \cup_{i=1}^3 C_i\} = 1 \text{ or equivalently } \mathbb{P}\{X \notin C_4\} = 0.$$

Lemma 3 Suppose (A1) and (A2) hold, and $X^h \Rightarrow X$. Then, $(X^h, \tau^h) \Rightarrow (X, \tau)$ as $h \rightarrow 0$.

PROOF: Let $\pi : D[0, 1] \mapsto [0, 1]$ be the mapping $\pi(x) = \inf\{t : x(t) \notin (\alpha(t), \beta(t))\}$, and $\hat{\pi} : D[0, 1] \rightarrow D[0, 1] \times [0, 1]$ defined by $\hat{\pi}(x) = (x, \pi(x))$. Note that we want to show that, under (A1) and (A2)

$$X^h \Rightarrow X \text{ implies that } \hat{\pi}(X^h) \Rightarrow \hat{\pi}(X).$$

Since (A2) implies that $\mathbb{P}\{X \in \cup_{i=1}^3 C_i\} = 1$, by the continuous mapping theorem [3, Theorem 2.7], the conclusion holds provided that the mapping $\hat{\pi}$ is almost surely continuous under the probability induced by $\mathbb{P}X^{-1}$, or equivalently $\hat{\pi}$ is continuous at arbitrary $x \in \cup_{i=1}^3 C_i$. Hence, it's enough to prove that π is continuous at arbitrary given $x \in \cup_{i=1}^3 C_i$ with respect to the Skorohod topology in the rest of the proof.

We assume $-\alpha(t) + \beta(t) < \infty$ for each t without loss of generality. Fix $x \in \cup_{i=1}^3 C_i$, and $\{x_n\}$ be an arbitrary sequence in $D[0, 1]$ satisfying $\|x_n - x\|_s \rightarrow 0$ as $n \rightarrow \infty$. By the definition of $\|\cdot\|_s$, it implies that there exists $\lambda_n \in \Lambda$ such that

$$\|\lambda_n - I\| \rightarrow 0, \quad \|x_n \circ \lambda_n - x\| \rightarrow 0, \text{ as } n \rightarrow \infty. \quad (6)$$

- (Case 1). Suppose $x \in C_1$, and define, for any $\varepsilon > 0$

$$\delta_\varepsilon := \frac{1}{2} \sup_{\pi(x) < t < \pi(x) + \varepsilon} (x(t) - \beta(t)).$$

By the definition of C_1 , we have $\delta_\varepsilon > 0$ for all $\varepsilon > 0$. Moreover, by continuity of x , there exists a time $t_\varepsilon \in (\pi(x), \pi(x) + \varepsilon)$ such that $x(t_\varepsilon) > \beta(t_\varepsilon) + \delta_\varepsilon$. Next, we observe that

$$|x_n(\lambda_n(t_\varepsilon)) - x(t_\varepsilon)| \leq \|x_n \circ \lambda_n - x\| \rightarrow 0, \text{ as } n \rightarrow \infty.$$

This implies that for some large $N_{\varepsilon,1}$

$$x_n(\lambda_n(t_\varepsilon)) \geq x(t_\varepsilon) - \frac{\delta_\varepsilon}{2} > \beta(t_\varepsilon) + \frac{\delta_\varepsilon}{2}, \quad \forall n \geq N_{\varepsilon,1}. \quad (7)$$

On the other hand, there exists $r_\varepsilon > 0$ due to the continuity of β such that

$$\sup_{t_\varepsilon - r_\varepsilon < t < t_\varepsilon + r_\varepsilon} |\beta(t) - \beta(t_\varepsilon)| < \frac{\delta_\varepsilon}{4}.$$

Moreover, since $\|\lambda_n - I\| \rightarrow 0$, we have for some large enough $N_{\varepsilon,2}$,

$$\|\lambda_n - I\| < r_\varepsilon, \quad \forall n \geq N_{\varepsilon,2}.$$

Therefore, it leads to

$$|\beta(\lambda_n(t_\varepsilon)) - \beta(t_\varepsilon)| \leq \sup_{t_\varepsilon - r_\varepsilon < t < t_\varepsilon + r_\varepsilon} |\beta(t) - \beta(t_\varepsilon)| < \frac{\delta_\varepsilon}{4}, \quad \forall n \geq N_{\varepsilon,2}. \quad (8)$$

(7) together with (8) implies

$$x_n(\lambda_n(t_\varepsilon)) \geq \beta(\lambda_n(t_\varepsilon)) + \frac{\delta_\varepsilon}{4}, \quad \forall n \geq N_\varepsilon := \max\{N_{\varepsilon,1}, N_{\varepsilon,2}\},$$

or equivalently,

$$\pi(x_n) \leq \lambda_n(t_\varepsilon) \leq \lambda_n(\pi(x) + \varepsilon), \quad \forall n \geq N_\varepsilon.$$

Finally, taking the $\limsup_{n \rightarrow \infty}$ on each side of the above inequality, and using $\|\lambda_n - I\| \rightarrow 0$ of (6), we have

$$\limsup_{n \rightarrow \infty} \pi(x_n) \leq \limsup_{n \rightarrow \infty} \lambda_n(\pi(x) + \varepsilon) = \pi(x) + \varepsilon.$$

So we conclude that by arbitrariness of ε

$$\limsup_{n \rightarrow \infty} \pi(x_n) \leq \pi(x).$$

- (Case 2). Again suppose $x \in C_1$, and we prove the reverse inequality. By the definition of the first hitting time, for any $\varepsilon > 0$, there exists $\delta_\varepsilon > 0$ such that

$$(\beta - x)^*(\pi(x) - \varepsilon) > \delta_\varepsilon, \quad (x - \alpha)^*(\pi(x) - \varepsilon) > \delta_\varepsilon. \quad (9)$$

Also, (6) implies that, $\|\alpha \circ \lambda_n - \alpha\| \rightarrow 0$, $\|\beta \circ \lambda_n - \beta\| \rightarrow 0$ and $\|x_n \circ \lambda_n - x\| \rightarrow 0$. Using the triangle inequality, we have

$$\|(x_n - \alpha) \circ \lambda_n - (x - \alpha)\| \rightarrow 0, \quad \|(x_n - \beta) \circ \lambda_n - (x - \beta)\| \rightarrow 0.$$

So, there exists a large N_ε such that

$$\|(x_n - \alpha) \circ \lambda_n - (x - \alpha)\| < \frac{\delta_\varepsilon}{2}, \quad \|(x_n - \beta) \circ \lambda_n - (x - \beta)\| < \frac{\delta_\varepsilon}{2}, \quad \forall n \geq N_\varepsilon.$$

In particular, this leads to

$$(\alpha - x_n) \circ \lambda_n(t) < (\alpha - x)(t) + \frac{\delta_\varepsilon}{2}, \quad (x_n - \beta) \circ \lambda_n(t) < (x - \beta)(t) + \frac{\delta_\varepsilon}{2}, \quad \forall t \in [0, 1], \quad n \geq N_\varepsilon.$$

By taking $\sup_{[0, \pi(x) - \varepsilon]}$ on both sides of the above inequalities and using (9), we have

$$(x_n - \beta)^* \lambda_n(\pi(x) - \varepsilon) < (x - \beta)^*(\pi(x) - \varepsilon) + \frac{\delta_\varepsilon}{2} < -\frac{\delta_\varepsilon}{2}, \quad \forall n \geq N_\varepsilon,$$

and

$$(\alpha - x_n)^* \lambda_n(\pi(x) - \varepsilon) < (\alpha - x)^*(\pi(x) - \varepsilon) + \frac{\delta_\varepsilon}{2} < -\frac{\delta_\varepsilon}{2}, \quad \forall n \geq N_\varepsilon.$$

Thus, we have

$$\pi(x_n) \geq \lambda_n(\pi(x) - \varepsilon), \quad \forall n \geq N_\varepsilon.$$

Taking $\liminf_n$ and using (6) and the fact of the arbitrariness of ε , we have

$$\liminf_n \pi(x_n) \geq \pi(x).$$

Summarizing the above, we have proved the continuity of π at $x \in C_1$ with respect to the Skorohod topology, i.e., $\lim_{n \rightarrow \infty} \pi(x_n) = \pi(x)$ whenever $x_n \rightarrow x \in C_1$ with respect to the Skorohod topology. Similar arguments as in Cases 1 and 2 yield that π is also continuous at $x \in C_2$. Moreover, π is continuous at $x \in C_3$ thanks to Lemma 2. This completes the proof. $\square$

Lemma 4 *Suppose (A1) and (A2) hold. If $(X^h, \tau^h) \Rightarrow (X, \tau)$ as $h \rightarrow 0$, then*

1. $(X^h, X^{h,*}, \tau^h) \Rightarrow (X, X^*, \tau)$ as $h \rightarrow 0$,
2. $(\Pi_{\mathcal{T}^h}(X^h), X^{h,*}(\tau^h), \tau^h) \Rightarrow (\Pi_{\mathcal{T}}(X), X^*(\tau), \tau)$ as $h \rightarrow 0$.

PROOF: By Lemma 2, the mapping $\pi_1^* : D[0, 1] \times [0, 1] \mapsto D^2[0, 1] \times \mathbb{R}$ by $\pi_1^*(x, t) = (x, x^*, t)$ is continuous in the product space in the Skorohod topology, where $D^2[0, 1] = D[0, 1] \times D[0, 1]$. Therefore, $(X^h, \tau^h) \Rightarrow (X, \tau)$ implies $\pi_1^*(X^h, \tau^h) \Rightarrow \pi_1^*(X, \tau)$ by mapping theorem, which in turn concludes the first result $(X^h, X^{h,*}, \tau^h) \Rightarrow (X, X^*, \tau)$.

Next, for simplicity, we only consider the special case with $(c_1, c_2) = (1/2, 1)$, so that the desired result is reduced to show that

$$(X^h(\tau^h), X^{h,*}(\tau^h), X^h(\tau^h/2), \tau^h) \Rightarrow (X(\tau), X^*(\tau), X(\tau/2), \tau), \quad \text{as } h \rightarrow 0.$$

Let $\bar{\pi} : D^2[0, 1] \times [0, 1] \mapsto \mathbb{R}^3$ by

$$\bar{\pi}(x, y, t) = (x(t), y(t), x(t/2)), \forall (x, y, t) \in D^2[0, 1] \times [0, 1].$$

Fix (x, y, t) such that (x, y) is continuous at t and $t/2$, respectively. Suppose $(x_n, y_n, t_n) \rightarrow (x, y, t)$ in their product space. Then, there exists $\lambda_n \in \Lambda$ such that

$$\|\lambda_n - I\| \rightarrow 0, \quad \|x_n - x \circ \lambda_n\| + \|y_n - y \circ \lambda_n\| + |t_n - t| \rightarrow 0, \quad \text{as } n \rightarrow \infty. \quad (10)$$

Therefore,

$$\begin{aligned} |x_n(t_n) - x(t)| &\leq |x_n(t_n) - x(\lambda_n(t_n))| + |x(\lambda_n(t_n)) - x(t)| \\ &\leq \|x_n - x \circ \lambda_n\| + |x(\lambda_n(t_n)) - x(t)| \end{aligned}$$

Note that the first term $\|x_n - x \circ \lambda_n\| \rightarrow 0$ as $n \rightarrow \infty$ by (10). On the other hand, since $t_n \rightarrow t$ and $\|\lambda_n - I\| \rightarrow 0$ as $n \rightarrow \infty$, we have $|\lambda_n(t_n) - t| \rightarrow 0$ as $n \rightarrow \infty$. Hence, the second term $|x(\lambda_n(t_n)) - x(t)| \rightarrow 0$ by continuity of x at t . So we have $x_n(t_n) \rightarrow x(t)$. Similarly we can show $y_n(t_n) \rightarrow y(t)$ and $x_n(t_n/2) \rightarrow x(t/2)$. Thus, we conclude

$\bar{\pi}$ is continuous at (x, y, t) whenever (x, y) is continuous at t and $t/2$.

Note that, (X, X^*) is continuous on $[0, 1]$ almost surely in $\mathbb{P}$. In particular, $X(\cdot, \omega)$ and $X^*(\cdot, \omega)$ are continuous at $\tau(\omega)$ and $\tau(\omega)/2$ almost surely in ω with respect to $\mathbb{P}$. Therefore, if we set

$$A := \{\omega \in \Omega : (X(\cdot, \omega), X^*(\cdot, \omega)) \text{ is continuous at } \tau(\omega) \text{ and } \tau(\omega)/2\},$$

then it's equivalent to $\mathbb{P}(A) = 1$. Therefore, with random element $Z : \Omega \mapsto (X, X^*, \tau)$, we have $\mathbb{P}Z^{-1}(D^2[0, 1] \times [0, 1]) = \mathbb{P}(A) = 1$, and $\bar{\pi}$ is continuous almost surely in $\mathbb{P}Z^{-1}$. This also implies the mapping $\hat{\pi} : D^2[0, 1] \times [0, 1] \mapsto \mathbb{R}^4$ by

$$\hat{\pi}(x, y, t) := (x(t), y(t), x(t/2), t) = (\pi(x, y, t), t), \forall (x, y, t) \in D^2[0, 1] \times [0, 1]$$

is continuous almost surely in $\mathbb{P}Z^{-1}$. Applying mapping theorem [3, Theorem 2.7], together with the previous result of $(X^h, X^{h,*}, \tau^h) \Rightarrow (X, X^*, \tau)$, we conclude the second result. $\square$

Next, we make an assumption on the payoff function g .

(A3) $g : \mathbb{R}^{m+2} \rightarrow \mathbb{R}$ is an almost surely continuous function with respect to $\mathbb{P}Z^{-1}$, where $Z := (\Pi_{\mathcal{T}}(X), X^*(\tau), \tau)$. Moreover, g satisfies either one of the following growth conditions:

1. g is a bounded function;
2. g is a function with linear growth and $\{X^h(t) : h > 0, t \in [0, 1]\}$ is uniformly integrable.

The following result is a consequence of Lemma 4.

Proposition 5 *Assume (A1)-(A3). Let X be a $\mathcal{F}_t$ -adapted continuous process with initial $X(0) = x$, and X^h be a sequence of RCLL processes satisfying $X^h \Rightarrow X$ as $h \rightarrow 0$. Then, $\lim_{h \rightarrow 0} V^h = V$.*

PROOF: Since (A1) and (A2) hold, Lemma 4 implies that

$$(\Pi_{\mathcal{T}^h}(X^h), X^{h,*}(\tau^h), \tau^h) \Rightarrow (\Pi_{\mathcal{T}}(X), X^*(\tau), \tau)$$

as $h \rightarrow 0$. Together with (A3), it results in $\lim_{h \rightarrow 0} V^h = V$; see [3, P. 25, 31]. $\square$

Now, we are ready to answer question (Q1).

Theorem 6 *Suppose (A1)-(A3) hold. If $X^h \Rightarrow X$ as $h \rightarrow 0$, then $\lim_{h \rightarrow 0} V^h = V$.*

PROOF: This is a consequence of Lemmas 3, 4, and Proposition 5. $\square$

Theorem 6 holds under assumptions of (A1)-(A3). Recall that (A1) is not a restriction. We will discuss (A2), (A3), and the uniform integrability in what follows.

Remark 7 (Discussions on (A2) and Example 1) In fact, (A2) is a requirement on the regularity of the boundary ∂Q with respect to the process X , and it is referred to as τ' -regularity for simplicity; see [17]. Note that since X in Example 1 violates τ' -regularity (A2), by observing

$$\inf\{t > \tau : X(t) \notin [\alpha(t), \beta(t)]\} \wedge 1 = 1 > 1/2 = \tau,$$

it yields the convergence to the wrong value. In other words, (A2) is crucial for the investigation of the convergence.

Remark 8 (Discussions on (A3) and Example 2) Assumption (A3) is the requirement on the function g . First of all, it allows discontinuity of g , but it can not be too much discontinuous in the sense that it is at least required to be almost surely continuous. However, it is already enough to include option pricing for the discontinuous payoff, such as barrier option.

In (A3), another issue yet mentioned is the growth condition of g . In particular, if g is of linearly growth function of the underlying price like in the call type option, then one shall verify uniform integrability. For instance, Example 2 converges to a wrong value since $\{X(t) : t \in (0, 1)\}$ is not uniformly integrable while the payoff function $g(x) = x$ is linear growth. Recall the definition of uniform integrability. A set of random variables $\{Y_\gamma : \gamma \in \Gamma\}$ is said uniformly integrable, if for any $\varepsilon > 0$, there exists a compact set K such that

$$\sup_{\gamma \in \Gamma} \mathbb{E}[Y_\gamma I_{\{Y_\gamma \notin K\}}] < \varepsilon.$$

Next, we present a proposition to ease the verification of the uniform integrability, which will be used later practically. The proof is referred to [7].

Proposition 9 *A family of real-valued random variable $\{Y_\gamma : \gamma \in \Gamma\}$ for some index set Γ is uniformly integrable if $\sup_{\gamma} \mathbb{E}[|Y_\gamma|^p] < \infty$ for some $p > 1$.*

3 Weak Convergence of Markov Chain Approximation

3.1 Markov Chain Approximation

In this section, we establish the weak convergence of Markov chain approximation for multidimensional stochastic differential equations. Let $Y = \{Y(t) : t \in [0, 1]\}$ be the unique weak solution of

$$dY(t) = b(Y(t), t)dt + \sigma(Y(t), t)dW(t); Y(0) = y, \quad (11)$$

where $b : \mathbb{R}^{d+1} \mapsto \mathbb{R}^d$, and W is a standard $\mathbb{R}^{d_1}$ Brownian motion, and $\sigma : \mathbb{R}^{d+1} \mapsto \mathbb{R}^{d \times d_1}$.

Let $t_0^h = 0 \leq t_1^h \leq \dots \leq t_N^h = 1$ be a sequence of increasing stopping times with respect to a discrete filtration $\{\mathcal{F}_i^h : i = 0, 1, \dots\}$, and $\{Y_i^h : i = 1, 2, \dots, N\}$ be a sequence $\mathcal{F}_i^h$ measurable Markov chain in $\mathbb{R}^d$ with transition probability

$$\mathbb{P}\{Y_{i+1}^h \in dy | Y_i^h = x, t_i^h = t\} = p^h(t, x, y).$$

We use $Y^h = \{Y^h(t) : t \in [0, 1]\}$ to denote piecewise constant interpolation

$$Y^h(t) = \sum_{i=0}^{n-1} Y_i^h I_{\{t_i^h \leq t < t_{i+1}^h\}}. \quad (12)$$

For notational simplicity, we set $\Delta t_n^h = t_{n+1}^h - t_n^h$ and $\Delta Y_n^h = Y_{n+1}^h - Y_n^h$. The interpolation of the Markov chain process Y^h is said to be *locally consistent*, if

$$(LC1) \quad \mathbb{E}[\sup_n \Delta t_n^h] = O(h),$$

$$(LC2) \quad \mathbb{E}[\Delta Y_n^h | Y_n^h = y, t_n^h = t] = \mathbb{E}[\Delta t_n^h | Y_n^h = y, t_n^h = t] \cdot (b(y, t) + O(h)),$$

$$(LC3) \quad \text{cov}(\Delta Y_n^h | Y_n^h = y, t_n^h = t) = \mathbb{E}[\Delta t_n^h | Y_n^h = y, t_n^h = t] \cdot ((\sigma\sigma')(y, t) + O(h)).$$

In local consistency (LC2) and (LC3), $O(h)$ is understood as a d -dimensional vector or $d \times d$ dimensional matrix with each element being $O(h)$, whenever it is appropriate.

(A4) b and σ are Lipschitz in y and Hölder-1/2 continuous in t , i.e., with $\phi = b, \sigma$,

$$|\phi(y_1, t_1) - \phi(y_2, t_2)| \leq K(|y_1 - y_2| + |t_1 - t_2|^{1/2}).$$

Theorem 10 *For the stochastic differential equation (11), assume that (A4) is satisfied, and Y^h of (12) is a sequence of random processes satisfying the local consistency condition. Then, the family of random variables $\{Y^h(t) : t \in [0, 1], h > 0\}$ are uniformly integrable, and $Y^h \Rightarrow Y$ as $h \rightarrow 0$.*

PROOF: We divide the proof into several steps.

1. First, we perform the following useful estimates. Set $\mathbb{E}_n^h[\cdot] := \mathbb{E}[\cdot | Y_n^h = y, t_n^h = t]$, and

$$z^h(t) := \max\{n \geq 0 : t_n^h \leq t\}, \quad \Delta M_n^h = \Delta Y_n^h - \mathbb{E}_n^h[\Delta Y_n^h].$$

Then

$$\begin{aligned} \mathbb{E}[\sup_{0 \leq n \leq z^h(t)-1} |\sum_{i=0}^n \Delta Y_i^h|^2] &= \mathbb{E}[\sup_{0 \leq n \leq z^h(t)-1} |\sum_{i=0}^n \mathbb{E}_i^h[\Delta Y_i^h] + \Delta M_i^h|^2] \\ &\leq K \mathbb{E}\left[\sup_{0 \leq n \leq z^h(t)-1} \left|\sum_{i=0}^n \mathbb{E}_i^h[\Delta Y_i^h]\right|^2\right] + K \mathbb{E}\left[\sup_{0 \leq n \leq z^h(t)-1} \left|\sum_{i=0}^n \Delta M_i^h\right|^2\right] := \tilde{I}_1 + \tilde{I}_2. \end{aligned}$$

We estimate the above two terms in the last line separately. We have

$$\begin{aligned}
\tilde{I}_1 &= \mathbb{E} \left[\sup_{0 \leq n \leq z^h(t)-1} \left| \sum_{i=0}^n \mathbb{E}_i^h [\Delta Y_i^h] \right|^2 \right] \\
&\leq \mathbb{E} \left[\sup_{0 \leq n \leq z^h(t)-1} \left(\sum_{i=0}^n \mathbb{E}_i^h [\Delta t_i^h] (|b|(Y_i^h, t_i^h) + O(h)) \right)^2 \right], \quad \text{by (LC2)} \\
&\leq \mathbb{E} \left[\left(\sum_{i=0}^{z^h(t)-1} \mathbb{E}_i^h [\Delta t_i^h] (|b|(Y_i^h, t_i^h) + O(h)) \right)^2 \right] \\
&\leq \mathbb{E} \left[\left(\sum_{i=0}^{z^h(t)-1} \Delta t_i^h (|b|(Y_i^h, t_i^h) + O(h)) \right)^2 \right], \quad \text{by Tower conditional expectation} \\
&\leq K \mathbb{E} \left[\left(\sum_{i=0}^{z^h(t)-1} \Delta t_i^h \right) \left(\sum_{i=0}^{z^h(t)-1} \Delta t_i^h |b|^2(Y_i^h, t_i^h) \right) \right] + K t^2 h^2, \\
&\leq K t \mathbb{E} \left[\sum_{i=0}^{z^h(t)-1} \Delta t_i^h |b|^2(Y_i^h, t_i^h) \right] + K t^2 h^2,
\end{aligned}$$

and

$$\begin{aligned}
\tilde{I}_2 &= \mathbb{E} \left[\sup_{0 \leq n \leq z^h(t)-1} \left| \sum_{i=0}^n \Delta M_i^h \right|^2 \right] \\
&\leq \mathbb{E} \left[\sum_{i=0}^{z^h(t)-1} |\Delta M_i^h|^2 \right], \quad \text{by Doob's Maximal inequality} \\
&= \mathbb{E} \left[\sum_{i=0}^{z^h(t)-1} |\Delta M_i^h|^2 \right], \quad \text{since } \mathbb{E}[(\Delta M_i^h)'(\Delta M_j^h)] = 0, \forall i \neq j \\
&\leq K \mathbb{E} \left[\sum_{i=0}^{z^h(t)-1} \Delta t_i^h \text{tr}(\sigma \sigma')(Y_i^h, t_i^h) + (\Delta t_i^h)^2 |b|^2(Y_i^h, t_i^h) \right] + K h t, \quad \text{by (LC2) (LC3),}
\end{aligned}$$

where $\text{tr}(A)$ denotes the trace of A . Therefore, we have

$$\begin{aligned}
&\mathbb{E} \left[\sup_{0 \leq n \leq z^h(t)-1} \left| \sum_{i=0}^n \Delta Y_i^h \right|^2 \right] \\
&\leq K(t+1) \mathbb{E} \left[\sum_{i=0}^{z^h(t)-1} \Delta t_i^h (|b|^2 + \text{tr}(\sigma \sigma'))(Y_i^h, t_i^h) \right] + K h t.
\end{aligned} \tag{13}$$

2. Applying inequality (13), we have

$$\begin{aligned}
\mathbb{E}[|Y^h(t)|^2] &\leq K \mathbb{E}[|y|^2 + \sum_{i=0}^{z^h(t)-1} \Delta Y_i^h]^2 \\
&\leq K|y|^2 + K \mathbb{E} \left[\sum_{i=0}^{z^h(t)-1} \Delta t_i^h (|b|^2 + \text{tr}(\sigma \sigma'))(Y_i^h, t_i^h) \right] + K h t \\
&\leq K|y|^2 + K \mathbb{E} \left[\int_0^t (|b|^2 + \text{tr}(\sigma \sigma'))(Y^h(s), s) ds \right] + K \\
&\leq K + K \mathbb{E} \left[\int_0^t (Y^h(s))^2 ds \right]
\end{aligned}$$

Gronwall's inequality then yields

$$\mathbb{E}[|Y^h(t)|^2] \leq K \text{ for all } t \in [0, T]. \quad (14)$$

Thus we conclude $\{Y^h(t) : 0 \leq t \leq T, h > 0\}$ is uniformly integrable by Proposition 9.

3. We consider the tightness of $\{\mathbb{P}_h = \mathbb{P}(Y^h)^{-1} : h > 0\}$. It is enough to verify conditions imposed on Theorem 13.3.2 and the subsequent corollary given in [3].

(a) By Chebyshev's inequality and (14)

$$\lim_{a \rightarrow \infty} \limsup_{h \rightarrow 0} \mathbb{P}\{|Y^h(t)| \geq a\} \leq \lim_{a \rightarrow \infty} \limsup_{h \rightarrow 0} \frac{1}{a^2} \mathbb{E}[|Y^h(t)|^2] = 0.$$

(b) For the purpose of characterization of tightness for discontinuous functions, we need to introduce some notions of modulus of continuity $\omega(Y^h, \delta)$ and $\omega'(Y^h, \delta)$ as the following. First, define $\omega(Y^h, \mathcal{I}) = \sup_{r_1, r_2 \in \mathcal{I}} |Y^h(r_1) - Y^h(r_2)|$ for any subset $\mathcal{I} \subset [0, 1]$. We also use $\mathcal{T}(\delta)$ to denote the collection of all δ -sparse partitions of $[0, 1]$. Then, we can define, for any $\delta > 0$

$$\omega(Y^h, \delta) = \sup_{0 \leq t \leq 1-\delta} \omega(Y^h, [t, t+\delta]),$$

and

$$\omega'(Y^h, \delta) = \inf_{\{t_i\} \in \mathcal{T}(\delta)} \max_i \omega(Y^h, [t_{i-1}, t_i]).$$

For the purpose of tightness of discontinuous functions, we also need to introduce modified version of modulus of continuity. For an arbitrary $\varepsilon > 0$, repeating the arguments of (13), we have

$$\begin{aligned} & \lim_{\delta \rightarrow 0} \limsup_{h \rightarrow 0} \mathbb{P}\{\omega'(Y^h, \delta) \geq \varepsilon\} \\ & \leq \lim_{\delta \rightarrow 0} \limsup_{h \rightarrow 0} \mathbb{P}\{\omega(Y^h, 2\delta) \geq \varepsilon\} \\ & \leq \lim_{\delta \rightarrow 0} \limsup_{h \rightarrow 0} \frac{1}{\varepsilon^2} \mathbb{E}[\omega^2(Y^h, 2\delta)] \\ & \leq \lim_{\delta \rightarrow 0} \limsup_{h \rightarrow 0} \frac{1}{\varepsilon^2} \mathbb{E}\left[\sup_{z^h(t)-1 \leq n \leq z^h(t+2\delta)-1} \left|\sum_{i=0}^n \Delta Y_i^h\right|^2\right] \\ & \leq \lim_{\delta \rightarrow 0} \limsup_{h \rightarrow 0} \frac{1}{\varepsilon^2} \mathbb{E}\left[\sup_{z^h(t)-1 \leq n \leq z^h(t+2\delta)-1} \sum_{i=0}^n \Delta t_i^h (|b|^2 + \text{tr}(\sigma\sigma'))(Y_i^h, t_i^h)\right] + Kh\delta \\ & \leq \lim_{\delta \rightarrow 0} \limsup_{h \rightarrow 0} \frac{1}{\varepsilon^2} \mathbb{E}\left[\int_t^{t+2\delta} K(|Y^h(s)|^2 + 1)ds\right] + Kh\delta \\ & \leq \lim_{\delta \rightarrow 0} \limsup_{h \rightarrow 0} K\delta \left(\frac{1}{\varepsilon^2} + h\right) = 0. \end{aligned}$$

As a result, $\{\mathbb{P}_h = \mathbb{P}(Y^h)^{-1} : h > 0\}$ is tight.

4. Since Y^h is tight, for an arbitrary infinite sequence, there exists a subsequence that has a weak limit. For notational convenience, we denote this subsequence again by $\{Y^h\}$, and its limit by $\bar{Y}$. Due to uniqueness of weak solution, it is enough to show that $\bar{Y}$ is the weak solution of (11). Since $Y^h \Rightarrow \bar{Y}$, the uniform integrability and (A4) lead to

$$\begin{aligned} \lim_{h \rightarrow 0} \mathbb{E}[Y^h(t)] &= \mathbb{E}[\bar{Y}(t)], \quad \text{and} \\ \lim_{h \rightarrow 0} \mathbb{E}[b(Y^h(t), t)] &= \mathbb{E}[b(\bar{Y}(t), t)]. \end{aligned}$$

Therefore, if we set

$$M(t) := \bar{Y}(t) - \bar{Y}(0) - \int_0^t b(\bar{Y}(s), s) ds,$$

we have

$$\begin{aligned} \mathbb{E}[M(t)] &= \lim_{h \rightarrow 0} \mathbb{E}[Y^h(t) - Y^h(0) - \int_0^t b(Y^h(s), s) ds] \\ &= \lim_{h \rightarrow 0} \mathbb{E} \left[\sum_{i=1}^{z^h(t)} ((Y_i^h - Y_{i-1}^h) - \int_{t_{i-1}^h}^{t_i^h} b(Y_{i-1}^h, t) dt) \right] - \lim_{h \rightarrow 0} \mathbb{E} \left[\int_{t_{z^h(t)}^h}^t b(Y_{z^h(t)}^h, s) ds \right] \end{aligned} \quad (15)$$

Note that regarding the last term in the above, by linear growth of b

$$\lim_{h \rightarrow 0} \mathbb{E} \left[\left| \int_{t_{z^h(t)}^h}^t b(Y_{z^h(t)}^h, s) ds \right| \right] \leq \lim_{h \rightarrow 0} \mathbb{E} \left[\int_{t_{z^h(t)}^h}^t K + K|Y_{z^h(t)}^h| ds \right] \leq K \lim_{h \rightarrow 0} \mathbb{E}[\Delta t_{z^h(t)}^h (1 + |Y^h(t)|)].$$

Since the Skorohod representation theorem (see [3, Page 70]) allows us to work with a common probability space $(\Omega_1, \mathcal{F}_1, \mathbb{P}_1)$ on which $(\Delta t_{z^h}^h, Y^h)$ converges to $(0, \bar{Y})$ almost surely. Together with uniform integrability, it leads to

$$\lim_{h \rightarrow 0} \mathbb{E}[\Delta t_{z^h(t)}^h (1 + |Y^h(t)|)] = \mathbb{E}[\lim_{h \rightarrow 0} \Delta t_{z^h(t)}^h (1 + |Y^h(t)|)] = 0.$$

Therefore, we conclude that the second term on (15) satisfies

$$\lim_{h \rightarrow 0} \mathbb{E} \left[\int_{t_{z^h(t)}^h}^t b(Y_{z^h(t)}^h, s) ds \right] = 0.$$

As for the first term of (15), using the Hölder continuity in t and the tower property on local consistency

$$\begin{aligned} &\lim_{h \rightarrow 0} \mathbb{E} \left[\sum_{i=1}^{z^h(t)} ((Y_i^h - Y_{i-1}^h) - \int_{t_{i-1}^h}^{t_i^h} b(Y_{i-1}^h, t) dt) \right] \\ &= \lim_{h \rightarrow 0} \mathbb{E} \left[\sum_{i=1}^{z^h(t)} ((Y_i^h - Y_{i-1}^h) - b(Y_{i-1}^h, t) \Delta t_{i-1}^h + O(|\Delta t_{i-1}^h|^{3/2})) \right] \\ &= \lim_{h \rightarrow 0} \mathbb{E} \left[\sum_{i=1}^{z^h(t)} (\Delta t_{i-1}^h O(h) + O(|\Delta t_{i-1}^h|^{3/2})) \right] = 0. \end{aligned}$$

Therefore, $\mathbb{E}[M(t)] = 0$. In fact, one uses exactly the same procedure to show $\mathbb{E}[M(t)|\mathcal{F}_s] = M(s)$ for any $0 \leq s \leq t$, and it concludes $M(t)$ is a martingale process.

Next, we use $\bar{Y}_l$ and Y_l^h to denote the l th component of the vector process $\bar{Y}$ and Y^h , and use $\langle \bar{Y}_l, \bar{Y}_m \rangle(t)$ to denote the cross-variation of two real processes Y_l and Y_m up to time t . Then,

$$\begin{aligned} & \mathbb{E}|\langle \bar{Y}_l, \bar{Y}_m \rangle(t) - \int_0^t (\sigma_l \sigma_m)(\bar{Y}(s), s) ds| \\ &= \lim_{h \rightarrow 0} \mathbb{E}|\langle Y_l^h, Y_m^h \rangle(t) - \int_0^t (\sigma_l \sigma_m)(Y^h(s), s) ds| \\ &= \lim_{h \rightarrow 0} \mathbb{E}|\sum_{i=1}^n (Y_{l,i}^h - Y_{l,i-1}^h)(Y_{m,i}^h - Y_{m,i-1}^h) - \int_0^t (\sigma_l \sigma_m)(Y^h(s), s) ds| = 0. \end{aligned}$$

The last equality can be obtained similar to the above owing to $\mathbb{E}[M(t)] = 0$, the local consistency, and regularity assumption (A4) on σ . Therefore, the cross-variation $\langle \bar{Y}_l, \bar{Y}_m \rangle(t) = \int_0^t (\sigma_l \sigma_m)(\bar{Y}(s), s) ds$ in L^1 and in probability. This again implies that $M(t)$ is a d -dimensional martingale process with its quadratic variation $\langle M \rangle(t) = \int_0^t (\sigma \sigma')(\bar{Y}(s), s) ds$ in probability. Applying Levy's martingale characterization on the time-changed Brownian motion, there exists a d -dimensional Brownian motion $B(t)$ such that $M(t) = \int_0^t \sigma(\bar{Y}(s), s) dB(s)$. Therefore, $\bar{Y}$ is the weak solution of (11).

Summarizing all the above, it leads to the conclusion. $\square$

3.2 Examples of Weak Convergence of Markov Chain Approximation

The above construction of Markov chain approximation is based on the local consistency, which generalizes the local consistency in [14, Theorem 10.4.1] in the following senses. First, σ and b may be unbounded but have linear growth. Therefore, the geometric Brownian motion is covered by weak convergence result of Theorem 10 as an important application. Secondly, from the definition of local consistency in (9.4.2) of [14], Δt_n^h has to be $\mathcal{F}_n^h$ measurable. However, the local consistency (LC1)-(LC3) allows us to have Δt_n^h to be only $\mathcal{F}_{n+1}^h$ measurable.

In fact, locally consistent MC approximation is flexible for its various choices. For illustrations, we give several simple MC approximations for one dimensional process different from [14], which can be easily verified to be locally consistent.

Example 3 Euler approximation can be considered as a special case of MC approximations. Let $\{Y_n^h\}$ be a Markov chain generated by

1. $Y_0^h = y, t_0^h = 0$.

2. Let the transition probability be $\Delta t_n^h = h$

$$Y_{n+1}^h = Y_n^h + b(Y_n^h, nh)h + \sigma(Y_n^h, nh)\sqrt{h}N_n, \quad (16)$$

where $\{N_n\}$ are i.i.d. standard normal random variables. One can easily verify all local consistency, and Theorem 10 implies, assuming (A4), the piecewise constant interpolation Y^h of the form (12) is convergent to Y of (11).

Example 4 The following MC approximation can be considered as an extension of binomial approximation of Brownian motion, since it is adapted by approximation of $W(t+h^2) - W(t) \sim \text{Bernoulli}(1/2)$. Let $d = d_1 = 1$. Let $\{Y_n^h\}$ be a Markov chain generated by

1. $Y_0^h = y, t_0^h = 0$.

2. Let the transition probability is, with $\Delta t_n^h = h^2$

$$\mathbb{P}\left(Y_{n+1}^h = Y_n^h + b(nh^2, Y_n^h)h^2 \pm \sigma(nh^2, Y_n^h)h \mid Y_n^h\right) = 1/2 \quad (17)$$

The above Markov chain is locally consistent since direct computation leads to

1. $\mathbb{E}[\sup_n \Delta t_n^h] = h^2$,

2. $\mathbb{E}[\Delta Y_n^h \mid Y_n^h = y, t_n^h = t] = \mathbb{E}[\Delta t_n^h \mid Y_n^h = y, t_n^h = t] \cdot b(y, t)$,

3. $\text{cov}(\Delta Y_n^h \mid Y_n^h = y, t_n^h = t) = \mathbb{E}[\Delta t_n^h \mid Y_n^h = y, t_n^h = t] \cdot \sigma^2(y, t)$.

Therefore, assuming (A4), the piecewise constant interpolation Y^h of the form (12) is convergent to Y of (11) by Theorem 10.

Example 5 This is a binomial tree type approximation of the diffusion term $\sigma(Y(t), t)dW(t)$ in (11). Let $d = d_1 = 1$. Let $\{Y_n^h\}$ be a Markov chain generated by

1. $Y_0^h = x, t_0^h = 0$.

2. Let $\Delta t_n^h(Y_n^h, t_n^h) = \frac{h^2}{\sigma^2(Y_n^h, t_n^h)} \wedge h^2$, and the transition probability is

$$\begin{aligned} \mathbb{P}\left(Y_{n+1}^h = Y_n^h + b(Y_n^h, t_n^h)\Delta t_n^h(Y_n^h, t_n^h) \pm h \mid (Y_n^h, t_n^h), \sigma^2(Y_n^h, t_n^h) \geq h\right) &= 1/2 \\ \mathbb{P}\left(Y_{n+1}^h = Y_n^h + b(Y_n^h, t_n^h)\Delta t_n^h(Y_n^h, t_n^h) \mid (Y_n^h, t_n^h), \sigma^2(Y_n^h, t_n^h) < h\right) &= 1. \end{aligned} \quad (18)$$

Note that, this MC is also locally consistent since

1. $\mathbb{E}[\sup_n \Delta t_n^h] \leq h^2$,

2. $\mathbb{E}[\Delta Y_n^h | Y_n^h = y, t_n^h = t] = \mathbb{E}[\Delta t_n^h | Y_n^h = y, t_n^h = t] \cdot b(y, t),$
3. $\text{cov}(\Delta Y_n^h | Y_n^h = y, t_n^h = t) = \mathbb{E}[\Delta t_n^h | Y_n^h = y, t_n^h = t] \cdot (\sigma^2(y, t) + O(h)).$

Therefore, assuming (A4), the piecewise constant interpolation Y^h of the form (12) is convergent to Y of (11) by Theorem 10.

3.3 Can We Expect the Strong Convergence?

In general, the answer is NO. To illustrate this point, we consider a special case of Euler approximation of Example 3. Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, on which $\mathcal{F}_t$ is filtration satisfying general conditions, and W_t of (11) is a standard 1-d Brownian motion. We construct strong approximation of Euler-Maruyama's method by taking on N_n of (16) by

$$N_n = \frac{W_{n+1} - W_n}{\sqrt{h}}.$$

Under assumption (A4), the SDE (11) has unique strong solution. Suppose $\hat{Y}^h$ is a continuous interpolation of Euler approximation of Y^h given by

$$\hat{Y}^h(t) = \hat{Y}_{nh}^h + b(nh, \hat{Y}_{nh}^h)(t - nh) + \sigma(nh, \hat{Y}_{nh}^h)(W(t) - W(nh)), \quad \text{for } t \in [nh, nh + h).$$

Then a classical result shows that see for example, [15, Theorem 2.7.3],

$$\mathbb{E} \left[\sup_{0 \leq t \leq T} |Y(t) - \hat{Y}^h(t)| \right] \leq Kh^{1/2}.$$

However, the above inequality fails for the piecewise constant interpolation of EM approximation $\{Y^h\}$. Otherwise, we have following simple counter example. Consider EM approximation of W_t on $[0, 1]$ by equal step size $h = 1/N$. Then, we have

$$\mathbb{E} \left[\sup_{0 \leq t \leq 1} |W(t) - W([Nt]/N)| \right] = \mathbb{E} \left[\sup_{1 \leq n \leq N} \sup_{(n-1)/N \leq t < n/N} |W(t) - W(\frac{n-1}{N})| \right].$$

Note that, $\bar{W}(t) = \sqrt{N}W(t/N)$ is a standard BM on time-scaled filtration. So one can reduce the above equality as

$$\mathbb{E} \left[\sup_{0 \leq t \leq 1} |W(t) - W([Nt]/N)| \right] = \frac{1}{\sqrt{N}} \mathbb{E} \left[\sup_{1 \leq n \leq N} \Lambda_n \right],$$

where $\{\Lambda_n\}$ are i.i.d. random variables defined by

$$\Lambda_n = \sup_{n-1 \leq t < n} |\bar{W}(t) - \bar{W}(n-1)|.$$

Since, Λ_n 's are unbounded iid random variables, $\mathbb{E} \left[\sup_{1 \leq n \leq N} \Lambda_n \right]$ goes to infinity as $N \rightarrow \infty$. This shows that

$$\mathbb{E} \left[\sup_{0 \leq t \leq 1} |W(t) - W([Nt]/N)| \right] > O(N^{-1/2}).$$

In conclusion, one cannot expect more than weak convergence merely under local consistency.

4 Ramification and Further Remarks

4.1 Application to Discretely Monitoring Barrier Option Underlying Stochastic Volatility Model

We begin this section with an application of Theorem 6 to the following stochastic volatility model; see [1]. Let W and B are two standard Brownian motions with correlation ρ in the filtered probability space $(\Omega, \mathcal{F}, \mathbb{P}, \mathcal{F}_t)$. Suppose the stock price follows

$$dX(t) = X(t)\sigma(Y(t))dW(t) \quad (19)$$

with initial $X(0) = x > 0$, and volatility factor follows

$$dY(t) = Y(t)(\mu(t)dt + b(t)dB(t)) \quad (20)$$

with initial $Y(0) = y > 0$. We consider a discretely monitoring barrier option price given by (3).

One can check that X and Y are unique nonnegative strong solution of SDEs (19) and (20), if σ satisfies polynomial growth, and μ and b are Hölder-1/2 continuous. In addition, we assume

$$\text{non-degeneracy of } X, \text{ i.e., } \sigma(y) > 0 \text{ for all } y > 0.$$

Note that it is possible to have $\sigma(0) = 0$ under the above assumption. We examine the convergence $\lim_h V^h = V$ when X^h is constructed by the Euler approximation.

Note that the regularity condition (A2) is satisfied by [16, Proposition A.1] due to $\sigma(\cdot) > 0$. On the other hand, the payoff function

$$g(x_1, x_2, \dots, x_m, x_{m+1}, t) = (x_m - \frac{1}{2})^+ I_{\{\max\{x_1, \dots, x_m\} > 1\}}$$

is only discontinuous at the points in the set $\{\max_i X(i/m) = 1\}$. Thanks to the fact

$$\mathbb{P}\{\max_i X(i/m) = 1\} = 0,$$

it implies that g is almost surely continuous with respect to $\mathbb{P}$. Another thing yet to be verified is the uniform integrability of $\{X^h(t) : h, t\}$, since g is linear growth. Thanks to Theorem 10 together with (A4), we have $X^h \Rightarrow X$ and the desired uniform integrability holds. Therefore, we reach the affirmative answer $\lim_h V^h = V$.

4.2 Further Remarks

This work has been devoted to analyzing approximation to path-dependent functionals. We have obtained weak convergence of the approximation schemes, which provide a unified approach for

proving the convergence of numerical approximation of path-dependent functionals for a wide range of applications.

This paper focused on diffusion models. For future work, it is worthwhile to examine systems driven by pure jump processes, jump diffusions, and systems with an additional factor process such as nowadays popular regime-switching processes. Systems with memory (time delays) form another class of important problems. Much work can also be devoted to numerical solutions of various stochastic differential equations, coordination of multi-agent systems, and many Monte Carlo optimization problems in which one needs to treat path-dependent functionals.

References

- [1] E. Bayraktar, K. Kardaras, and H. Xing. Valuation equations for stochastic volatility models. *SIAM Journal on Financial Mathematics*, 3:351–373, 2012.
- [2] E. Bayraktar, Q.S. Song, and J. Yang. On the continuity of stochastic control problems on bounded domains. *Stochastic Analysis and Applications*, 29(1):48–60, 2011.
- [3] P. Billingsley. *Convergence of probability measures*. Wiley Series in Probability and Statistics: Probability and Statistics. John Wiley & Sons Inc., New York, second edition, 1999. A Wiley-Interscience Publication.
- [4] M. Broadie, P. Glasserman, and S. G. Kou. Connecting discrete and continuous path-dependent options. *Finance Stoch.*, 3:55–82, 1999.
- [5] B. de Saporta, F. Dufour, K. Gonzalez, Numerical method for optimal stopping of piecewise deterministic Markov processes. *Ann. Appl. Probab.*, Vol. 20, pp. 1607-1637, (2010).
- [6] F. Dufour and A.B. Piunovskiy, Multiobjective stopping problem for discrete-time Markov processes: convex analytic approach, *J. Appl. Probab.*, Vol. 47, pp. 947-966, (2010).
- [7] R. Durrett. *Probability*. The Wadsworth & Brooks/Cole Statistics/Probability Series. Wadsworth & Brooks/Cole Advanced Books & Software, Pacific Grove, CA, 3rd edition, 2005. Theory and examples.
- [8] P. Glasserman. *Monte Carlo Methods In Financial Engineering*. Springer, 2004.
- [9] M. Jeanblanc, M. Yor, and M. Chesney. *Mathematical methods for financial markets*. Springer Finance. Springer-Verlag London Ltd., London, 2009.
- [10] I. Karatzas and S.E. Shreve. *Brownian motion and stochastic calculus, Graduate Texts in Mathematics*, volume 113. Springer-Verlag, New York, second edition, 1991.

- [11] P.E. Kloeden and E. Platen. *Numerical solution of stochastic differential equations*, volume 23 of *Applications of Mathematics (New York)*. Springer-Verlag, Berlin, 1992.
- [12] S. G. Kou. On pricing of discrete barrier options. *Statistica Sinica*, 13:955–964, 2003.
- [13] H.J. Kushner. Numerical methods for stochastic control problems in continuous time. *SIAM J. Control Optim.*, 28(5):999–1048, 1990.
- [14] H.J. Kushner and P. Dupuis. *Numerical methods for stochastic control problems in continuous time*, volume 24 of *Applications of Mathematics (New York)*. Springer-Verlag, New York, second edition, 2001.
- [15] X.R. Mao. *Stochastic Differential Equations and Applications*. Horwood Pub Ltd, 2007.
- [16] Q.S. Song, G. Yin, and C. Zhu. Optimal switching with constraints and utility maximization of an indivisible market. *SIAM Journal on Control and Optimization*, 50(2):629–651, 2012.
- [17] D. Stroock and S.R.S. Varadhan. On degenerate elliptic-parabolic operators of second order and their associated diffusions. *Comm. Pure Appl. Math.*, 25:651–713, 1972.
- [18] L. Szpruch and D.J. Higham, Comparing hitting time behavior of Markov jump processes and their diffusion approximations, *SIAM Multiscale Model. Simul.*, Vol. 8, pp. 605–621, (2010).